

AMENDMENT UNDER 37 CFR § 1.111  
Application No. 09/903,476

*[Signature]* Clean copy of claim 6

6. (Amended) The process of claim 5 wherein contacting the oxygen treated catalyst with the aqueous medium is conducted in the liquid phase.

Clean copy of claim 9

9. (Amended) The process of claim 5 wherein the molecular sieve of the alkylation catalyst is MCM-22, PSH-3, SSZ-25, MCM-36, MCM-49, MCM-56, faujasite, mordenite or zeolite beta.

*[Signature]* Clean copy of claim 10

10. (Amended) The process of claim 5 wherein said aqueous medium is ammonium nitrate solution, ammonium carbonate solution or acetic acid solution.

*A2* *[Signature]* Clean copy of claim 11

11. (Amended) The process of claim 5 wherein contacting the catalyst with the aqueous medium is conducted at a temperature of about 15 to about 120° C for a period of about 10 minutes to about 48 hours.

*[Signature]* Clean copy of claim 12

12. (Amended) The process of claim 5 further including calcining the aqueous medium contacted catalyst at a temperature of about 25 to about 600° C for a period of about 10 minutes to about 48 hours.

Please add the following claims 13 to 24:

*Sub C2 > 1* 13. A process for alkylating an aromatic compound comprising:  
contacting an alkylatable aromatic compound and an alkylating agent with an alkylation catalyst comprising a molecular sieve under alkylation conditions; and  
when said alkylation catalyst has become at least partially deactivated, contacting said alkylation catalyst with an oxygen-containing gas at a temperature of about 120 to about 600° C; and then

2

AMENDMENT UNDER 37 CFR § 1.111  
Application No. 09/903,476

B  
B  
contacting the oxygen treated catalyst with an aqueous medium selected from ammonium nitrate solution or ammonium carbonate solution.

*the going up  
writing of*

14. The process of claim 13 wherein contacting the oxygen treated catalyst with the aqueous medium is conducted in the liquid phase.

3 15. The process of claim 13 wherein the alkylating agent includes an alkylating aliphatic group having 1 to 5 carbon atoms.

A 16. The process of claim 13 wherein the alkylating agent is ethylene or propylene and the alkylatable aromatic compound is benzene.

A 17. The process of claim 13 wherein the molecular sieve of the alkylation catalyst is MCM-22, PSH-3, SSZ-25, MCM-36, MCM-49, MCM-56, faujasite, mordenite or zeolite beta.

6 18. The process of claim 13 further including calcining the aqueous medium contacted catalyst at a temperature of about 25 to about 600° C for a period of about 10 minutes to about 48 hours.

5vb 3 > 7 19. A process for alkylating an aromatic compound comprising:  
contacting an alkylatable aromatic compound and an alkylating agent with an alkylation catalyst comprising a molecular sieve under alkylation conditions; and  
when said alkylation catalyst has become at least partially deactivated, contacting said alkylation catalyst with an oxygen-containing gas at a temperature of about 120 to about 600° C; and then

inner B 1  
contacting the oxygen treated catalyst with an aqueous medium, wherein the molecular sieve of the alkylation catalyst is PSH-3, SSZ-25, MCM-36, MCM-49, MCM-56, faujasite, mordenite or zeolite beta.

AMENDMENT UNDER 37 CFR § 1.111  
Application No. 09/903,476

8 20. The process of claim 19 wherein contacting the oxygen treated catalyst with the aqueous medium is conducted in the liquid phase.

9 21. The process of claim 19 wherein the alkylating agent includes an alkylating aliphatic group having 1 to 5 carbon atoms.

10 22. The process of claim 19 wherein the alkylating agent is ethylene or propylene and the alkylatable aromatic compound is benzene.

11 23. The process of claim 19 further including calcining the aqueous medium contacted catalyst at a temperature of about 25 to about 600° C for a period of about 10 minutes to about 48 hours.

24. The process of claim 19 wherein the aqueous medium is ammonium nitrate solution, ammonium carbonate solution or acetic acid solution.